ACCESS

Conference Abstract

The use of avian museum specimen data in phenology studies: prospects and challenges

Keith Barker[‡], Michael Wells[‡], Dakota Rowsey[‡]

PROCEEDINGS OF

‡ University of Minnesota, St. Paul, United States of America

Corresponding author: Keith Barker (barke042@umn.edu)

Received: 15 Aug 2017 | Published: 15 Aug 2017

Citation: Barker K, Wells M, Rowsey D (2017) The use of avian museum specimen data in phenology studies: prospects and challenges. Proceedings of TDWG 1: e20324. <u>https://doi.org/10.3897/tdwgproceedings.1.20324</u>

Abstract

Museum specimens offer a rich source of data on both long-term averages and temporal trends in organismal phenology. To date much of this work has focused on plants, but animal specimens are also useful in this regard. In particular, bird specimens may include data on age, gonad size and development, and molt, all of which are relevant to estimating breeding phenology. In addition, bird collections include nests and eggs, which are direct records of breeding. These data are some of the richest available for any vertebrate group, and the potential for integration with citizen-science-based data (e.g. eBird) is extremely promising. However, there are a number of challenges associated with use of these data, including biased geographic sampling, incomplete digitization, and non-uniform data standards, that currently limit their utility. In addition, methods for analysis of these data have not been well developed. Here, we outline informatics challenges associated with developing a database of avian gonad size data. In addition, we present a novel analytical framework for estimating phenology from such data (Fig. 1) that is broadly applicable to seasonal phenotype measurements.

© Barker K et al. This is an open access article distributed under the terms of the Creative Commons Attribution License (CC BY 4.0), which permits unrestricted use, distribution, and reproduction in any medium, provided the original author and source are credited.




Figure 1.

Phenology of *Molothrus* testis development, estimated using log-normal periodic regression (data from Payne 1973, *Condor* 75: 80-99).

Keywords

phenology, birds, gonad, periodic regression

Presenting author

Keith Barker

Funding program

Division of Environmental Biology, Systematics and Biodiversity Science Cluster

Grant title

DEB-1541312